

ADA035228

NAMRL-1230

13

Jr

NORMAL HEMATOLOGICAL, BIOCHEMICAL, AND SERUM ELECTROLYTE  
VALUES FOR A COLONY OF RHESUS MONKEYS (Macaca mulatta)

Lieutenant Colonel James L. Kupper, USAF VC,  
Captain Matthew J. Kessler, USAF VC, and Larry L. Cook

*[Handwritten signature]*



DDC  
RECEIVED  
FEB 7 1977  
A

28 October 1976

Approved for public release; distribution unlimited.

Approved for public release; distribution unlimited.

NORMAL HEMATOLOGICAL, BIOCHEMICAL, AND SERUM ELECTROLYTE  
VALUES FOR A COLONY OF RHESUS MONKEYS (Macaca mulatta)

Lieutenant Colonel James L. Kupper, USAF VC,  
Captain Matthew J. Kessler, USAF VC, and Larry L. Cook

Naval Medical Research and Development Command  
XSB09.ED6.6-B1

Approved by

Ashton Graybiel, M.D.  
Assistant for Scientific Programs

Released by

Captain R.E. Mitchel, MC USN  
Commanding Officer

28 October 1976

NAVAL AEROSPACE MEDICAL RESEARCH LABORATORY  
PENSACOLA, FLORIDA 32508

## SUMMARY PAGE\*

### THE PROBLEM


To establish normal hematological, biochemical, and serum electrolyte values for rhesus monkeys housed at the Naval Aerospace Medical Research Laboratory.

### FINDINGS

Venipuncture was performed on 56 male and 57 female rhesus monkeys (*Macaca mulatta*) according to a standardized protocol to determine normal hematological, biochemical, and serum electrolyte values. Arithmetic means and standard deviations are tabulated for each parameter for each sex. The results from previously reported studies on rhesus monkeys are also tabulated and, where possible, values among the studies are compared.

### ACKNOWLEDGMENTS

The services provided by NAMRL Biomedical Sciences Division and Veterinary Sciences Division personnel are appreciated. The authors are especially grateful to Don Personette, Alvin L. Armstrong, and HM<sub>2</sub> J.D. Clayton for their diligent efforts and assistance in the preparation of this report.



\*The animals used in this study were handled in accordance with the Principles of Laboratory Animal Care established by the Committee on the Guide for Laboratory Animal Resources, National Academy of Science-National Research Council.

## INTRODUCTION

A number of reports have been published in the scientific literature which detail the normal hematological, biochemical, and serum electrolyte values for captive rhesus monkeys (1-6, 8-22, 24-28). Information available from those studies was found to be of limited value for our purposes for various reasons. Sampling techniques, the use of pharmacological agents, handling procedures, and variations in laboratory methods and equipment all contributed to the differences reported among normal data and the results routinely obtained at the Naval Aerospace Medical Research Laboratory.

The specific purpose of this report is to establish normal values for our colony that were obtained under controlled and reproducible conditions to aid in the selection of monkeys for use in future experimentation.

## SUBJECTS AND PROCEDURE

The subjects involved were rhesus monkeys purchased as a single shipment of 120 animals from another military research laboratory. The monkeys were maintained for 10 months prior to the study on a commercial laboratory primate chow\* that was supplemented regularly with fresh apples and oranges. The sampled population consisted of 56 and 57 female monkeys whose ages were determined by dentition (7). The males ranged in age from 24 months to 54 months, with an arithmetic mean of 42.2 months and a standard deviation of 5.8 months. Female monkeys ranged from 24 months to 80 months, with an arithmetic mean of 50.1 months and a standard deviation of 8.8 months.

The male monkeys ranged in weight from 3.38 to 8.45 kilograms, with an arithmetic mean of 5.01 kilograms and a standard deviation of 1.2 kilograms. The females ranged in weight from 3.13 kilograms to 6.99 kilograms, with an arithmetic mean of 4.67 kilograms and a standard deviation of 0.74 kilograms.

The monkeys all received a comprehensive clinical physical examination. Of the 120 animals maintained in the colony, 113 monkeys were found to be healthy and in good physical condition prior to sampling. Among the seven monkeys not available for testing were those with major disabilities (amputated limbs) or severe illnesses (salmonellosis), and those that had died prior to the sampling date.

The subjects were deprived of all food for 17 hours prior to venipuncture, but were allowed water ad lib. Thirty minutes prior to

---

\*Purina Monkey Chow<sup>®</sup>, Ralston Purina Co., St. Louis, Missouri

venipuncture, the monkeys were administered a neuroleptic-narcotic combination drug, fentanyl-droperidol\*, at a dosage of 1 cc per 40 pounds of body weight (1 cc per 18.2 kilograms of body weight) intramuscularly (23). This dosage regimen was sufficient to produce moderate catalepsis in the subjects and facilitate venipuncture. The animals were restrained in a horizontal position. The vast majority of blood samples were obtained from the femoral veins but, occasionally, samples were inadvertently obtained from the femoral arteries. Ten cubic centimeters of blood were collected using dry, sterile, disposable, 12-cc plastic syringes and 20-gauge, 1-inch disposable needles. For routine blood counts the needle was removed from the tip of the syringe and 1 cc of blood was transferred without delay into a glass test tube that contained 1 drop of heparin at a concentration of 10,000 I.U. per cubic centimeter. The remaining 9 cc of whole blood were transferred from the syringe into glass test tubes that contained no anticoagulant. Blood smears were made from a drop of the heparinized blood sample.

#### METHODS

1. Hematology - Total erythrocyte count, total white blood cell count, and hematocrit were determined by using the Model ZBI Coulter counter. The counter was equipped with a channelizer which displayed the cell-size distributions for accurate setting of the upper and lower window of the counter. Hemoglobin concentration was determined using a Coulter Electronics, Inc., hemoglobinometer. The procedure involved was the cyanomethemoglobin-colometric method. Mean corpuscular value (MCV), mean corpuscular hemoglobin (MCH), and mean corpuscular hemoglobin concentration (MCHC) were calculated from data obtained from the Coulter counter and Coulter hemoglobinometer. Differential white blood cell counts were performed by counting 100 white blood cells. An experienced hematology technician familiar with rhesus monkeys performed the hematology studies.

2. Biochemistry - The Gilford Model 3500 automated chemistry analyzer was used for the measurement of serum chemistries in this study. Prepared reagents were obtained from Worthington Biochemical Corporation for the following chemistries: total protein, albumin, serum glutamic pyruvic transaminase, serum glutamic oxaloacetic transaminase, lactate dehydrogenase, creatinine phosphokinase, glucose, and triglycerides. Abbott Laboratories supplied the prepared reagents for blood urea nitrogen and cholesterol measurements. Serum lipoproteins and proteins were determined electrophoretically.

a. Total serum protein - The biuret reaction was employed in which cupric ions react with proteins in alkaline solution to form a blue-violet colored complex. A total protein standard of known protein concentration was used as a reference in the Gilford 3500.

---

\*Innovar , McNeil Laboratories, Fort Washington, PA

b. Serum albumin - Albumin reacts with the anionic bromcresol green dye and causes an increase in the green color. This color is measured at 628 nm and is directly proportional to the albumin concentration as compared to a serum albumin standard.

c. Serum glutamic oxaloacetic transaminase (SGOT) - The activity of SGOT is determined by coupling the reaction it catalyzes to a secondary reaction. In this secondary reaction NADH is oxidized to NAD, permitting the transaminase activity to be determined by the rate of decrease in absorbance at 340 nm.

d. Serum glutamic pyruvic transaminase (SGPT) - The activity of SGPT is proportional to the rate of a secondary reaction in which NADH is oxidized to NAD. This rate is measured spectrophotometrically at 340 nm.

e. Lactate dehydrogenase (LDH) - Lactate dehydrogenase catalyzes the conversion of lactate to pyruvate and NAD to NADH. LDH activity is directly proportional to the increase in absorbance of NADH measured at 340 nm.

f. Creatine phosphokinase (CPK) - CPK catalyzes the conversion of creatine phosphate and ADP to creatine and ATP. The ATP produced and glucose are converted to ADP and glucose-6-phosphate by hexokinase in a secondary reaction. In a tertiary reaction catalyzed by glucose-6-phosphate dehydrogenase, NAD is reduced to NADH. The rate of NADH production measured at 340 nm is proportional to the CPK activity.

g. Serum glucose - In the primary reaction hexokinase catalyzes the phosphorylation of glucose to glucose-6-phosphate. Glucose-6-phosphate is oxidized and NAD reduced in the secondary reaction by glucose-6-phosphate dehydrogenase. Since both reactions are essentially irreversible, the concentration of NADH produced in the secondary reaction, measured at 340 nm, is a direct measurement of the total glucose present in the sample serum when compared to a glucose standard.

h. Triglycerides - Serum triglycerides are first saponified with ethanolic potassium hydroxide to produce glycerol and free fatty acids. The free fatty acids are precipitated with magnesium sulfate. The glycerol is then measured by three coupled enzymatic reactions catalyzed by glycerol kinase, phosphokinase, and lactate dehydrogenase. In the third reaction NADH is oxidized to NAD. The total NADH decrease, measured at 340 nm, is directly proportional to the concentration of glycerol present after the saponification process and, therefore, to the concentration of triglycerides present in the sample.

i. Blood urea nitrogen (BUN) - Urease splits urea into ammonia and carbon dioxide. Glutamic dehydrogenase catalyzes the combination of the ammonia quantitatively with alpha-ketoglutarate along with the oxidation of NADH to NAD. The resultant decrease in absorbance at 340 nm is directly proportional to the concentration of ammonia formed which,

in turn, is quantitatively related to the concentration of urea initially present. This procedure is based on the comparison of the unknown samples to BUN standards.

j. Cholesterol - Cholesterol esters in serum are hydrolyzed to free cholesterol by cholesterol ester hydrolase. The free cholesterol produced is oxidized by cholesterol oxidase to cholest-4-en-3-one and hydrogen peroxide. The hydrogen peroxide couples with 4-aminoantipyrine and phenol in the presence of peroxidase to yield a quinoneimine dye with an absorption maximum at 500 nm. The amount of color produced is directly proportional to the total cholesterol concentration of the sample as compared to cholesterol standards.

k. Lipoprotein - Lipoproteins were determined by electrophoresis on Gelman Sephrophore III cellulose acetate membranes, using a Beckman Microzone Cell. The membranes were stained using the Kohn ozonization method and measured with a Beckman Model CUS-100 computing densitometer.

l. Protein electrophoresis - The process was the same as that described for lipoprotein electrophoresis except that oil Red O was used for staining the Gelman Sephrophore III membranes.

### 3. Serum Electrolytes

a. Calcium - A manual fluorescent titration method was used to determine serum calcium concentration. The dye calcein fluoresces in the presence of calcium. This fluorescent complex is titrated with ethylenediamine tetra-acetic acid (EDTA), the end point being the termination of fluorescence as observed under a longwave ultraviolet light. Reagents were obtained from Oxford Laboratories.

b. Sodium and potassium - These two electrolytes were determined by using the International Laboratory Flame Photometer Model 153.

## RESULTS AND DISCUSSION

Table I presents a summary of hematological values obtained from 56 male and 57 female monkeys sedated with Innovar<sup>®</sup>. Arithmetic means and standard deviations are tabulated for each sex group in Tables I-III.

The results of the biochemical serum analyses from the 113 sedated monkeys in the sampled population are listed in Table II, and the serum electrolyte values obtained from the same sample population of sedated monkeys are detailed in Table III.

Tables IV, V, and VI summarize the mean values of hematological, biochemical, and electrolyte values available in the literature on rhesus monkeys. Some authors reported results by male and female populations of monkeys, while others only reported values for both sexes combined. This resulted in some apparent discrepancies in Tables III, IV, and V. For instance, in Table IV, MCH determination, the range of means reported for combined sexes is greater than that for either sex group reported

separately. References are cited for each determination listed.

In general, the results accumulated from our subjects were consistent with previous findings reported in the literature. Notable exceptions include the following:

1. Hemoglobin values in grams percent were higher for the monkeys examined in the present study than in any other article reviewed. There is no obvious explanation for this inconsistency.

2. The female monkey mean corpuscular hemoglobin (MCH) in picograms was higher than all other reported values except for those of Robinson and Ziegler (19). The value reported for the male monkeys in our study exceeded all previous values reported in the literature.

3. The mean total white blood cell counts per cubic millimeter for our monkeys were generally lower than the means reported in most of the studies reviewed. The mean value of band neutrophils was higher than any of the other reported values. This is probably due to different interpretations of band cell types among hematology technicians.

4. The mean cholesterol values observed in our colony were lower than most values reported in the literature reviewed.

5. Potassium levels in millequivalents per liter for our female monkeys were below the normal range of reported mean values.

Normal mean values for lipoproteins (alpha, pre-beta, and beta), albumin, and globulins (alpha, beta, gamma) as a percentage of the total serum protein were not readily available in the literature. One study (22) reported serum lipoprotein profiles of twelve different primate species, but we were unable to compare those results with our values due to differences in analytical procedures and reporting units. Values for the normal mean triglyceride level in rhesus monkeys were also unavailable in published reports.

Values for normal serum glutamic oxaloacetic transaminase, serum glutamic pyruvic transaminase, lactate dehydrogenase, and creatine phosphokinase expressed in international units per liter were not found in other reports. Since these determinations are method specific, they cannot be compared with values obtained by using different procedures. A review of these parameters expressed in various units can be found in Bourne (4).



## REFERENCES

1. Anderson, D.R., Normal values for clinical blood chemistry tests of the Macaca mulatta monkey. Am. J. Vet. Res., 27: 1484-1489, 1966.
2. Back, K.C., Toxicity studies on animals exposed continuously to a 5 PSIA 100% oxygen environment for periods up to 235 days. In: Proceedings of the Second Annual Conference on Atmospheric Contamination in Confined Spaces. AMRL-TR-66-120. Wright-Patterson Air Force Base, Ohio: Aerospace Medical Research Laboratories, 1966. Pp 80-87.
3. Banerjee, S., and Chakrabarty, A.S., Anaemia and its relation with iron metabolism in scobotic monkeys. Indian J. Med. Res., 53: 835-848, 1965.
4. Bourne, G.H. (Ed.), The Rhesus Monkey, Volume II: Management, Reproduction, and Pathology. New York: Academic Press, 1975. Chapter 15.
5. Dillingham, L.A., Morrow, A.C., and Bronsdon, M.A., A comparison of the hemograms of Macaca mulatta, Macaca nemestrina and Papio anubis. Folia Primat., 14: 241-251, 1971.
6. Greep, R.O., Knobil, E., Hofmann, F.G., and Jones, T.L., Adrenal cortical insufficiency in the rhesus monkey. Endocrinology, 50: 664-676, 1952.
7. Hartmen, C.G., and Straus, W.L., jr. (Ed.), The Anatomy of the Rhesus Monkey (Macaca mulatta). New York: Hafner Publishing Co., 1933. P 21.
8. Huser, H.J., Atlas of Comparative Primate Hematology. New York: Academic Press, 1970. Pp 10 and 98.
9. Ives, M., and Dack, G.M., Alarm reaction and normal blood picture in Macaca mulatta. J. Lab. Clin. Med., 47: 723-729, 1956
10. Jones, E.S., McCall, K.B., Elvehjam, C.A., and Clark P.F., The effect of diet on the hemoglobin, erthrocytes, and leukocyte content of the blood of the rhesus monkey (Macaca mulatta). Blood, 2: 154-163, 1974.
11. King, T.O. and Gargus, J.L., Normal blood values of the adult female monkey (Macaca mulatta). Lab. Anim. Care, 17: 391-396, 1967.

12. Krise, G.M., Jr., and Wald, N., Normal blood picture of the Macaca mulatta monkey. J. Appl. Physiol., 12: 482-484, 1958.
13. MacEwan, J.D., and McNerney, J.M., Effect of a mixed gas atmosphere at 5 PSIA on the toxicity of NO<sub>2</sub> and O<sub>3</sub> in animals. In: Proceedings of Second Annual Conference on Atmospheric Contamination in Confined Spaces. AMRL-TR-66-120. Wright-Patterson Air Force Base, Ohio: Aerospace Medical Research Laboratories, 1966. Pp 290-295.
14. Martin, D.P., McGowan, M.J., and Loeb, W.F., Related changes of hematologic values in infant Macaca mulatta. Lab. Anim. Sci., 23: 194-200, 1973.
15. Melville, G.S., Whitcomb, W.H., and Martinez, R.S., Hematology of the Macaca mulatta monkey. Lab. Anim. Care, 17: 189-198, 1967.
16. Petery, J. J., Ultramicroanalysis of selected blood components of normal Macaca Mulatta. Lab. Anim. Care, 17: 342-344, 1967.
17. Pickering, D.E., and Kao, T.T.H., Fluid and electrolyte therapy for monkeys. J. Am. Vet. Med. Assoc., 138: 527-531, 1961.
18. Rao, G.N., and Shipley, E.G., Data on selected clinical blood chemistry tests of adult female rhesus monkeys (Macaca mulatta). Lab. Anim. Care, 20: 226-231, 1970.
19. Robinson, F.R., and Ziegler, R.F., Clinical laboratory data derived from 102 Macaca mulatta. Lab. Anim. Care, 18: 50-57, 1968.
20. Rollins, J.B., et al. Hematologic studies of the rhesus monkey (Macaca mulatta). Lab. Anim. Care, 20: 681-685, 1970.
21. Stanley, R.E., and Cramer, M.B., Hematologic values of the monkey (Macaca mulatta). Am. J. Vet. Res., 29: 1041-1047, 1968.
22. Srinivasan, S.R., McBride, J.R., Radhakrishnamurthy, B., and Berenson, G.S., Comparative studies on serum lipoprotein and lipid profiles in subhuman primates. Comp. Biochem. Physiol., 47B: 711-716, 1974.
23. Stunkhard, J.A., and Miller, J.C., An outline guide to general anesthesia in exotic species. Vet. Med. Small Anim. Clinician, Sept.: 1181-1186, 1974.
24. Switzer, J.W., Valerio, D.A., Martin, D.P., Valerjo, M.G., and Rininger, B.F., Hematologic changes associated with pregnancy and parturition in Macaca mulatta. Lab. Anim. Care, 20: 930-939, 1970.

25. Turbyfill, C.L., Cramer, M.B., Dewes, W.A., and Huguley, J.W., III, Serum and cerebral spinal fluid chemistry values for the monkey (Macaca mulatta). Lab. Anim. Care, 20: 269-273, 1970.
26. Van den Berghe, L., and Blitstein, I., Hematologic studies in the monkey. I. Cytologic composition of the blood. Ann. Soc. Belge. Med. Trop., 33: 709-729, 1953.
27. Vogin, E.E., and Oser, F., Comparative blood values in several species of nonhuman primates. Lab. Anim. Sci., 21: 937-941, 1971.
28. Vondruska, J.F., and Greco, R.A., Certain hematologic and blood chemical values in juvenile rhesus monkeys (Macaca mulatta). Bull. Amer. Soc. Vet. Clin. Pathol., 3: 27-36, 1974.

TABLE I  
Hematological Values for the NAMRL Rhesus Monkey Colony

	Males (N=56)		Females (N=57)	
	Mean	S.D.	Mean	S.D.
erythrocytes x $10^6/\text{mm}^3$	5.25	0.46	5.00	0.39
hemoglobin gm%	15.1	1.1	14.6	1.0
hematocrit %	42.9	4.4	41.8	4.0
MCV microns <sup>3</sup>	79.6	5.6	80.4	5.7
MCH picograms	29.4	1.1	26.7	1.4
MCHC %	38.3	2.4	32.8	1.8
leukocytes x $10^3/\text{mm}^3$	7.6	5.2	8.3	5.7
lymphocytes/100 WBC	43.8	9.6	57.3	9.5
neutrophils/100 WBC	32.0	8.4	35.2	8.6
band cells/100 WBC	1.8	1.4	1.9	1.0
monocytes/100 WBC	0.56	0.79	0.59	0.87
eosinophils/100 WBC	2.3	2.2	2.2	1.8
basophils/100 WBC	0.49	0.73	0.39	0.71

TABLE II  
Biochemical Values for the NAMRL Rhesus Monkey Colony

	Males (N=56)		Females (N=57)	
	Mean	S.D.	Mean	S.D.
total protein gm%	7.4	0.58	7.6	0.71
albumin gm%	4.2	0.38	4.0	0.40
cholesterol mg%	146.9	28.5	140.9	25.0
BUN mg%	23.5	4.4	22.8	4.7
glucose mg%	102.4	20.1	89.8	18.3
triglyceride mg%	57.7	19.8	56.2	12.1
SGOT I.U./liter	28.4	9.0	28.1	8.1
SGPT I.U./liter	25.9	8.0	24.4	8.7
LDH I.U./liter	348.1	214.7	364.0	177.3
CPK I.U./liter	159.8	133.3	164.7	95.0
albumin % total protein	61.5	4.7	60.5	6.2
globulins % total protein				
alpha	8.8	2.0	9.2	2.6
beta	17.9	3.3	17.0	3.7
gamma	11.0	4.4	13.3	5.2
lipoproteins % total lipo-				
proteins				
alpha	64.1	3.0	64.3	1.0
pre-beta	10.6	6.9	10.1	7.7
beta	26.4	7.5	26.6	7.4

TABLE III

## Serum Electrolyte Values for the NAMRL Rhesus Monkey Colony

mEq/L	Male(N=56)		Female(N=57)	
	Mean	S.D.	Mean	S.D.
calcium meq/liter	5.4	0.21	5.3	0.22
sodium meq/liter	147.2	3.8	147.5	5.1
potassium meq/liter	3.6	0.34	3.3	0.25

TABLE IV  
Summary of Reported Mean Hematological Values for Rhesus Monkeys

Determination	Combined Sexes	Males	Females	References
erythrocytes $\times 10^6/\text{mm}^3$	4.32-6.38	5.54-5.64	5.10-5.67	3, 4, 8-12, 15, 19, 21, 24, 26-28
hemoglobin gm%	11.7-13.5	13.2-12.3	12.3-13.4	3-5, 8-12, 15, 19-21, 24, 27, 28
hematocrit %	36.0-45.5	39.3-41.3	38.7-40.9	3-5, 8, 11, 12, 14, 15, 19, 20, 21, 24, 27, 28
MCV microns <sup>3</sup>	69.9-91.5	69.9-74.8	70.2-76.3	3, 4, 8, 15, 19, 21, 28
MCH picograms	22.0-28.1	23.7	23.7-24.3	3, 4, 8, 15, 19, 21, 28
MCHC %	28.9-35.1	32.0-34.4	31.9-34.2	3, 4, 8, 15, 19, 21, 28
leukocytes $\times 10^3/\text{mm}^3$	7.8-16.9	8.2-10.6	8.9-12.2	3-5, 8-12, 14, 15, 19-21, 24, 26-28
lymphocytes/100 WBC	45.3-72.8	56.3-59.6	55.1-57.9	4, 5, 8-12, 14, 15, 19, 21, 26-28
neutrophils/100 WBC	22.4-49.5	35.6-37.6	36.9-41.0	4, 5, 8-12, 14, 15, 19, 21, 26-28
bands/100 WBC	0-0.4	0	0	4, 5, 8-12, 14, 15, 19, 21, 26-28
monocytes/100 WBC	0-3.9	0-2.2	0-2.0	4, 5, 8-12, 14, 15, 19, 21, 26-28
eosinophils/100 WBC	0-5.2	0-2.3	0-1.7	4, 5, 8-12, 14, 15, 19, 21, 26-28
basophils/100 WBC	0-0.5	0-0.3	0-0.1	4, 5, 8-12, 14, 15, 19, 21, 26-28

TABLE V  
Summary of Reported Mean Biochemical Values for Rhesus Monkeys

Determination	Both Sexes	Males	Females	References
Total protein gm%	6.3-8.0	7.1-8.0	7.4-8.0	1,2,4,13,16,19,21,25,27
albumin gm%	3.6-5.0	4.2-5.0	4.4-4.7	1,2,4,6,13,19,20,25
cholesterol mg%	128-219	176	182	1,4,16,18,25,27
blood urea nitrogen mg%	10.1-25.3	10.1-22.2	12.8-22.9	1,4,16,18,20,25,27
glucose mg%	61.8-102.0	95.0-102.0	96.5-101.2	1,4,16-18,25,27



TABLE VI  
Summary of Reported Mean Electrolyte Values for Rhesus Monkeys

Determination	Both Sexes	Male	Female	References
calcium meq/liter	4.9-5.8	5.3	4.9	2,4,11,13,16,17,19,25
sodium meq/liter	139.1-153.0	155.1-157.3	154.0-156.1	1,2,4,6,13,17,19,25,27
potassium meq/liter	3.4-5.2	5.1-5.2	5.1	1,2,4,6,13,17,19,25,27

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NAMRL-1230	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Normal Hematological, Biochemical, and Serum Electrolyte Values for a Colony of Rhesus Monkeys, (Macaca mulatta)	5. TYPE OF REPORT & PERIOD COVERED	
7. AUTHOR(s) <del>1st Lieutenant Colonel</del> James L. Kupper, USAF VC, Colonel Matthew J. Kessler, USAF VC, and Larry L. Cook	6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Aerospace Medical Research Laboratory Pensacola, Florida 32508	8. CONTRACT OR GRANT NUMBER(s)	
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Medical Research and Development Command National Naval Medical Center Bethesda, Maryland 20034	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS  XSB09.ED6.6-B1.2	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	12. REPORT DATE 28 October 1976	
(12) 18p.	13. NUMBER OF PAGES 16	
	15. SECURITY CLASS. (of this report) Unclassified	
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) <u>Macaca mulatta</u> , Rhesus monkey, Blood values, Veterinary Medicine, Laboratory Animals		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) To establish normal hematological, biochemical, and serum electrolyte values for rhesus monkeys housed at the Naval Aerospace Medical Research Labora- tory, venipuncture was performed on 56 male and 57 female rhesus monkeys ( <u>Macaca</u> <u>mulatta</u> ). Arithmetic means and standard deviations are reported for each para- meter for sex. The results are compared with previously reported studies on rhesus monkeys from other laboratories.		

DD FORM 1473  
1 JAN 75EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-014-6001


UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

406061

**UNCLASSIFIED**

**SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)**



**UNCLASSIFIED**

**SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)**

